

Weyl Invariant Spacetime

Yutaka Hosotani

Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

Abstract

A new class of Weyl invariant backgrounds are presented in terms of the metric $G_{\mu\nu}$ and the anti-symmetric Kalb-Ramond fields $B_{\mu\nu}$. The ten-dimensional spacetime is a product of four-dimensional flat spacetime and curved six-dimensional spacetime having nonvanishing Ricci tensors. The non-vanishing Kalb-Ramond field strengths cannot be written globally as $H = dB$, being of the monopole type. Nevertheless they define homogeneous spacetime with no singularity.

1. Introduction

In the string theory the background spacetime cannot be arbitrarily assigned. For the consistency of the theory the Weyl invariance has to be maintained. At low energies the spacetime is described in terms of the metric $G_{\mu\nu}$, the anti-symmetric Kalb-Ramond fields $B_{\mu\nu}$, and the dilaton field Φ . In the background the trace of world-sheet energy momentum tensors is cast in the form

$$T^a{}_a = -\frac{1}{2\alpha'}\beta_{\mu\nu}^G g^{ab}\partial_a X^\mu\partial_b X^\nu - \frac{i}{2\alpha'}\beta_{\mu\nu}^B \epsilon^{ab}\partial_a X^\mu\partial_b X^\nu - \frac{1}{2}\beta^\Phi R \quad , \quad (1.1)$$

which must vanish to maintain the Weyl invariance. Here $1/(2\pi\alpha')$ and g_{ab} are the string tension and the world-sheet metric. ϵ^{ab} is the two-dimensional antisymmetric tensor. R is the scalar curvature of the spacetime. Callan et al. [1] have shown that the condition for having vanishing beta functions in the critical dimension $D = D_c$ leads to

$$\begin{aligned} \beta_{\mu\nu}^G &= \alpha' \left\{ (Ricci)_{\mu\nu} + 2\nabla_\mu \nabla_\nu \Phi - \frac{1}{4} H_{\mu\rho\sigma} H_\nu{}^{\rho\sigma} \right\} + O(\alpha'^2) = 0 \quad , \\ \beta_{\mu\nu}^B &= \alpha' \left\{ -\frac{1}{2} \nabla^\rho H_{\rho\mu\nu} + \nabla^\rho \Phi \cdot H_{\rho\mu\nu} \right\} + O(\alpha'^2) = 0 \quad , \\ \beta^\Phi &= \alpha' \left\{ -\frac{1}{2} \nabla^2 \Phi + \nabla_\mu \Phi \nabla^\mu \Phi - \frac{1}{24} H_{\mu\nu\lambda} H^{\mu\nu\lambda} \right\} + O(\alpha'^2) = 0 \quad . \end{aligned} \quad (1.2)$$

This set of the equations can be derived from the action

$$\begin{aligned} I &= \frac{1}{2\kappa^2} \int d^D x \, |\det G|^{1/2} \mathcal{L} \\ \mathcal{L} &= e^{-2\Phi} \left\{ R - \frac{1}{12} H_{\mu\nu\lambda} H^{\mu\nu\lambda} + 4\nabla_\mu \Phi \nabla^\mu \Phi + O(\alpha') \right\} \end{aligned} \quad (1.3)$$

by taking variation over $G_{\mu\nu}$, $B_{\mu\nu}$ and Φ . In other words Weyl invariant backgrounds are classical solutions to the equations of motion in the field theory defined by (1.3).

Most of the backgrounds considered in the literature are Ricci flat, $(Ricci)_{\mu\nu} = 0$, and have vanishing $H_{\mu\nu\lambda}$'s. It is of great interest to know if there are other kinds of backgrounds with nonvanishing $(Ricci)_{\mu\nu}$ and $H_{\mu\nu\lambda}$. [2] Here we loosen the conditions for the consistency. In the string theory it is assumed that the extra six dimensions are all spacelike, i.e. the signature of the metric is all positive. We drop this restriction.

We ask if there are new type of solutions to (1.2). Spacetime satisfying (1.2) is said, in the present paper, to be Weyl invariant.

At the bottom of this investigation is the observation that the Kalb-Ramond fields $B_{\mu\nu}$ are conformally invariant at the classical level and have natural configurations in six dimensions.[3] It is a remarkable fact that the string theory produces the Kalb-Ramond fields which may account for why we can see only four dimensions at very low energies. We shall see below that in one of the examples extra dimensions are compact and there appears high symmetry between the spacelike and timelike dimensions.

2. Consistency conditions

We shall impose another constraint to set up the analysis. We restrict ourselves to configurations with a constant dilaton field $\Phi = \text{constant}$. It seems very difficult to have configurations with spatially varying Φ when extra-dimensional spacetime is compact.

With this restriction imposed the consistency conditions (1.2) reduce, to $O(\alpha')$, to

$$\begin{aligned} (Ricci)_{\mu\nu} &= \frac{1}{4} H_{\mu\rho\sigma} H_{\nu}{}^{\rho\sigma} , \\ \nabla_{\rho} H^{\rho\mu\nu} &= \frac{1}{\sqrt{|G|}} \partial_{\rho} \left\{ \sqrt{|G|} H^{\rho\mu\nu} \right\} = 0 , \\ R &= 0 \quad , \quad H_{\mu\nu\lambda} H^{\mu\nu\lambda} = 0 . \end{aligned} \tag{2.1}$$

The last equality implies that, if the extra dimensions are all spacelike, i.e. the signature of the metric is all positive, all components of $H_{\mu\nu\lambda}$'s vanish so that the resultant space is Ricci flat. We shall consider more general spacetime.

The metric of six dimensional spacetime is written as

$$ds^2 = \eta_{ab} e^a \otimes e^b \tag{2.2}$$

where e^a 's ($a = 1 \sim 6$) are tetrad 1-forms. η_{ab} is a diagonal matrix, whose elements are either +1 or -1. In all of the Weyl invariant spacetime discussed bellow $\det(\eta_{ab})$ is -1. The spin connection 1-forms are defined by $de^a + \omega^a{}_b \wedge e^b = 0$. The curvature 2-forms are then given by $\mathcal{R}_{ab} = d\omega_{ab} + \omega_a{}^c \wedge \omega_{cb}$, whose components are Riemann tensors; $\mathcal{R}_{ab} = \frac{1}{2} R_{abcd} e^c \wedge e^d$. Ricci tensors are $(Ricci)_{ab} = R_{acb}{}^c$.

The Kalb-Ramond fields define 2-form fields; $B = \frac{1}{2}B_{\mu\nu} dx^\mu \wedge dx^\nu = \frac{1}{2}B_{ab} e^a \wedge e^b$. Their field strengths define 3-form fields $H = \frac{1}{6}H_{abc} e^a \wedge e^b \wedge e^c = dB$. The equations in (2.1) are rewritten as

$$(Ricci)_{ab} = \frac{1}{4} H_{acd} H_b{}^{cd} \equiv S_{ab} \quad , \quad (2.3)$$

$$R = 0 \quad , \quad (2.4)$$

$$dH = 0 \quad , \quad *d(*H) = 0 \quad . \quad (2.5)$$

$*$ denotes Hodge dual. We note that the B -fields are gauge fields for closed strings. The theory is invariant under a gauge transformation $B \rightarrow B + d\Lambda$ where Λ is a 1-form gauge function. Just like the monopole fields in the electrodynamics in three spatial dimensions, the potential B need not be defined globally. The field strengths H must satisfy (2.5).

3. Example I. $SU(2) \otimes \overline{SU(2)}$

A nontrivial Weyl-invariant background is obtained by considering a product of two three-spheres (S^3). Let us first recall some properties of a manifold S^3 . S^3 is isomorphic to a group manifold $SU(2)$. The relationship is most easily established by the correspondence

$$S^3 : y_1^2 + y_2^2 + y_3^2 + y_4^2 = 1 \Leftrightarrow \Omega = y_4 + i\vec{y} \cdot \vec{\tau} \in SU(2) \quad (3.1)$$

where τ^j 's ($j = 1, 2, 3$) are Pauli matrices. Maurer-Cartan 1-forms of $SU(2)$, $\sigma^j = -i\text{Tr} \tau^j \Omega^{-1} d\Omega$ satisfy $d\sigma^j = \frac{1}{2}\epsilon^{jkl}\sigma^k \wedge \sigma^l$. In terms of σ^j 's the metric of S^3 or $SU(2)$ is given by $ds^2 = \sum_{j=1}^3 \sigma^j \otimes \sigma^j$.

Now we prepare two sets of $SU(2)$;

$$d\sigma^j = \frac{1}{2}\epsilon^{jkl}\sigma^k \wedge \sigma^l \quad , \quad d\rho^j = \frac{1}{2}\epsilon^{jkl}\rho^k \wedge \rho^l \quad . \quad (3.2)$$

The metric of the six-dimensional manifold, $SU(2) \otimes \overline{SU(2)}$, is written as

$$ds^2 = \sum_{j=1}^3 (\sigma^j \otimes \sigma^j - \rho^j \otimes \rho^j) \quad , \quad (3.3)$$

or equivalently,

$$e^1 = \sigma^1, \quad e^2 = \sigma^2, \quad e^3 = \sigma^3, \quad e^4 = \rho^1, \quad e^5 = \rho^2, \quad e^6 = \rho^3, \\ \eta_{ab} = \text{diag}(1, 1, 1, -1, -1, -1) \quad . \quad (3.4)$$

Curvature 2-forms are given by

$$\mathcal{R}_{j,k} = \frac{1}{4} e_j \wedge e_k \\ \mathcal{R}_{j+3,k+3} = -\frac{1}{4} e_{j+3} \wedge e_{k+3} \\ \mathcal{R}_{j,k+3} = 0 \quad , \quad (j, k = 1, 2, 3) \quad . \quad (3.5)$$

Ricci tensors are found to be

$$(\text{Ricci})_{ab} = \frac{1}{2} \delta_{ab} = \frac{1}{2} \text{diag}(1, 1, 1, 1, 1, 1) \quad . \quad (3.6)$$

The scalar curvature $R = (\text{Ricci})_a^a$ vanishes. The length scale, L , in the metric has been set to be unity. Every quantity scales with L with an appropriate power.

We need to find 3-form field strengths H satisfying (2.3) and (2.5). In this connection it is very fruitful to recall how special configurations in non-Abelian gauge theory in four dimensions arise. They might have been realized in the early universe. If the universe is described by the Robertson-Walker metric with $k = +1$, namely with a spatial section S^3 , then there appears a natural mapping in the configurations of $SU(2)$ gauge fields. The 1-form gauge potential takes the form $A = f(t) \vec{\sigma} \cdot \vec{\tau}$ where σ^k 's are the Maurer-Cartan forms. This ansatz simultaneously solves the Einstein equations and Yang-Mills equations.[4, 5] The resultant space is homogeneous and isotropic. Its generalization solves the equations in the Einstein-Weinberg-Salam theory as well. The homogeneity of the space is maintained, but the space necessarily becomes anisotropic due to the $U(1)$ gauge interaction.[6] A crucial point is that 1-form gauge potentials have natural configurations on $SU(2) \sim S^3$.

In the problem under discussions we have the Abelian antisymmetric tensor fields B , namely 2-form fields. To our surprise there are natural configurations on the manifold (3.3). First notice that the 3-form field strength $H = d(\sigma^j \wedge \rho^k)$ is self-dual, $*H = +H$,

as straightforward calculations show. It therefore automatically solves $d(*H) = dH = 0$.

The symmetrized version

$$B_0 = \sum_{j=1}^3 \sum_{k=1}^3 \sigma^j \wedge \rho^k \quad , \quad H_0 = dB_0 \quad (3.7)$$

yields

$$\begin{aligned} dH_0 &= 0 \quad , \quad *d(*H_0) = 0 \quad , \\ (S_0)_{ab} &= \frac{1}{4} (H_0)_{acd} (H_0)_b{}^{cd} = -\frac{3}{2} \delta_{ab} \quad . \end{aligned} \quad (3.8)$$

The configuration H_0 solves almost all of the consistency equations except it gives a wrong sign for S_{ab} .

We next consider the following self-dual or anti-self-dual configuration;

$$\begin{aligned} H_{\pm} &= \sigma^1 \wedge \sigma^2 \wedge \sigma^3 \mp \rho^1 \wedge \rho^2 \wedge \rho^3 \\ &= e^1 \wedge e^2 \wedge e^3 \mp e^4 \wedge e^5 \wedge e^6 \\ *H_{\pm} &= \pm H_{\pm} \quad . \end{aligned} \quad (3.9)$$

It follows from (3.2) that $dH_{\pm} = 0$. We have

$$\begin{aligned} dH_{\pm} &= 0 \quad , \quad *d(*H_{\pm}) = 0 \quad , \\ (S_{\pm})_{ab} &= \frac{1}{4} (H_{\pm})_{acd} (H_{\pm})_b{}^{cd} = \frac{1}{2} \delta_{ab} \quad . \end{aligned} \quad (3.10)$$

$H_{\pm}$ itself solves (2.3) and (2.5). The configuration $H_{\pm}$ cannot be written globally as $H_{\pm} = dB_{\pm}$. Nevertheless it is a legitimate physical configuration as it solves the equations of motion (2.5). Its coupling to matter, say, strings, is well defined thanks to the gauge invariance associated with B . It is like a monopole configuration in the four-dimensional electromagnetism.[7] The vector potential for a magnetic monopole cannot be written globally without introducing a Dirac string.[8] We remark that the field strength $H_{\pm}$ is regular everywhere. It gives homogeneous spacetime.

General solutions to (2.3) and (2.5) in the metric (3.3) are

$$(i) \quad H = \pm H_+$$

$$(ii) \quad H = \alpha H_- + \beta H_0 \quad , \quad \alpha^2 - 3\beta^2 = 1 \quad . \quad (3.11)$$

As H_0 is self-dual, the interference term occurring in S_{ab} for the case (ii) vanishes. (3.3) and (3.11) define Weyl invariant spacetime.

The (anti-)self-duality relation for H has played a vital role. In six dimensions self-dual or anti-self-dual configurations are possible only when there are an odd number of timelike dimensions.

4. Example II. $SL(2, R) \otimes \overline{SL(2, R)}$

The second example of Weyl invariant spacetime is obtained by replacing $SU(2)$ by $SL(2, R)$ in the previous example. Generators of $SL(2, R)$ satisfy

$$[T_1, T_2] = T_3 \quad , \quad [T_2, T_3] = T_1 \quad , \quad [T_3, T_1] = -T_2 \quad . \quad (4.1)$$

The corresponding Maurer-Cartan 1-forms satisfy

$$d\sigma^1 = \sigma^2 \wedge \sigma^3 \quad , \quad d\sigma^2 = -\sigma^3 \wedge \sigma^1 \quad , \quad d\sigma^3 = \sigma^1 \wedge \sigma^2 \quad . \quad (4.2)$$

The metric of the manifold $SL(2, R)$ is given by $ds^2 = \sigma^1 \otimes \sigma^1 - \sigma^2 \otimes \sigma^2 + \sigma^3 \otimes \sigma^3$.

We prepare two sets of $SL(2, R)$ whose Maurer-Cartan 1-forms are denoted by $\{\sigma^j\}$ and $\{\rho^j\}$. The metric $ds^2 = \eta_{ab} e^a \otimes e^b$ of the six-dimensional manifold, $SL(2, R) \otimes \overline{SL(2, R)}$, is given by $e^j = \sigma^j$, $e^{j+3} = \rho^j$ ($j = 1, 2, 3$) and

$$\eta_{ab} = \text{diag} (1, -1, 1, -1, 1, -1) \quad . \quad (4.3)$$

Curvature 2-forms are given by

$$\begin{aligned} \mathcal{R}_{j,k} &= -\frac{1}{4} e_j \wedge e_k \\ \mathcal{R}_{j+3,k+3} &= +\frac{1}{4} e_{j+3} \wedge e_{k+3} \\ \mathcal{R}_{j,k+3} &= 0 \quad , \quad (j, k = 1, 2, 3) \quad . \end{aligned} \quad (4.4)$$

Compared with (3.5) in the previous example, the sign is opposite. Ricci tensors become

$$(Ricci)_{ab} = -\frac{1}{2} \text{diag} (1, -1, 1, 1, -1, 1) \quad . \quad (4.5)$$

The scalar curvature R vanishes.

The configurations solving (2.3) and (2.5) are

$$\begin{aligned}
H &= \pm H_{\pm} \ , \\
H_{\pm} &= \sigma^1 \wedge \sigma^2 \wedge \sigma^3 \pm \rho^1 \wedge \rho^2 \wedge \rho^3 \\
&= e^1 \wedge e^2 \wedge e^3 \pm e^4 \wedge e^5 \wedge e^6 \\
{}^*H_{\pm} &= \pm H_{\pm} \ .
\end{aligned} \tag{4.6}$$

Again these H 's cannot be written globally as $H = dB$. S_{ab} for these configurations is exactly given by the right hand side of (4.5), solving (2.3).

The gauge potential $B^{(jk)} = \sigma^j \wedge \rho^k$ yields anti-self-dual field strength; ${}^*(dB^{(jk)}) = -dB^{(jk)}$. However, a configuration of the type $B_0 = \sum_{j=1}^3 \sum_{k=1}^3 B^{(jk)}$ does not solve (2.3). On the manifold $SL(2, R) \otimes \overline{SL(2, R)}$ the allowed H configurations are (4.6) only. We remark that the manifold $SO(2, 2)$ is isomorphic to $SL(2, R) \otimes \overline{SL(2, R)}$.

5. Example III. $SU(2) \otimes SL(2, R)$ and $\overline{SU(2)} \otimes \overline{SL(2, R)}$

The third example is given by a product of $SU(2)$ and $SL(2, R)$. Let $\{\sigma^j\}$ and $\{\rho^j\}$ be Maurer-Cartan forms of $SU(2)$ and $SL(2, R)$, respectively. The metric is given by

$$ds^2 = \sigma^1 \otimes \sigma^1 + \sigma^2 \otimes \sigma^2 + \sigma^3 \otimes \sigma^3 + \rho^1 \otimes \rho^1 - \rho^2 \otimes \rho^2 + \rho^3 \otimes \rho^3 \ , \tag{5.1}$$

or equivalently $e^j = \sigma^j$, $e^{j+3} = \rho^j$ ($j = 1, 2, 3$) and $\eta_{ab} = \text{diag}(1, 1, 1, 1, -1, 1)$. The curvature 2-forms are given by (3.5) so that

$$(\text{Ricci})_{ab} = \frac{1}{2} \text{diag}(1, 1, 1, -1, 1, -1) \ , \tag{5.2}$$

yielding a vanishing scalar curvature. The Kalb-Ramond field strengths are

$$H = H_{\pm} = e^1 \wedge e^2 \wedge e^3 \mp e^4 \wedge e^5 \wedge e^6 = \pm {}^*H_{\pm} \ . \tag{5.3}$$

Eq. (2.3) is satisfied as

$$S_{ab} = \frac{1}{2} \text{diag}(1, 1, 1, -1, 1, -1) \ . \tag{5.4}$$

This time again $B^{(jk)} = \sigma^j \wedge \rho^k$ yields anti-self-dual field strengths; $*(dB^{(jk)}) = -dB^{(jk)}$.

The fourth example is obtained by flipping the signature of the third example, namely by $\overline{SU(2)} \otimes \overline{SL(2, R)}$. The curvature 2-forms are given by (4.4), whereas the Ricci tensors turn out the same as (5.2). The Kalb-Ramond field strengths must be $H = H_{\pm} = e^1 \wedge e^2 \wedge e^3 \pm e^4 \wedge e^5 \wedge e^6$. $B^{(jk)} = \sigma^j \wedge \rho^k$ yields self-dual field strengths on the manifold; $*(dB^{(jk)}) = +dB^{(jk)}$.

6. Other group manifolds: $SO(4)$ and $SO(3, 1)$

It may be worth clarifying why the manifolds $SO(4)$ and $SO(3, 1)$ are not Weyl invariant. The manifold $SO(4)$ is locally isomorphic to $SU(2) \otimes SU(2)$. The metric is given, in terms of the σ^j 's and ρ^j 's in Section 3, by $ds^2 = \sum_{j=1}^3 (\sigma^j \otimes \sigma^j + \rho^j \otimes \rho^j)$ instead of (3.3), which leads to (3.6) and nonvanishing scalar curvature $R = 3$. Hence the condition (2.4) cannot be satisfied. A varying dilaton field $\Phi(x)$ may give a nontrivial solution to (1.2). It, however, is difficult to have such Φ on a compact manifold.

The situation is similar for the manifold $SO(3, 1)$. This manifold is of special interest. If the ten-dimensional spacetime is a product of four-dimensional Minkowski spacetime and six-dimensional $SO(3, 1)$, then Lorentz symmetry transformations of the four-dimensional Minkowski spacetime have one-to-one correspondence to points or translations in the extra six-dimensional spacetime. It could be the origin of the local Lorentz invariance of the four-dimensional spacetime, when the four-dimensional spacetime is promoted to curved spacetime.

The manifold $SO(3, 1)$ is isomorphic to $SL(2, C)$ whose complex Maurer-Cartan 1-forms are denoted by $\sigma^k + i\rho^k$ ($k = 1, 2, 3$). They satisfy

$$d(\sigma^j + i\rho^j) = \frac{1}{2}\epsilon^{jkl}(\sigma^k + i\rho^k) \wedge (\sigma^l + i\rho^l) \quad . \quad (6.1)$$

The metric is

$$ds^2 = \sum_{j=1}^3 (\sigma^j \otimes \sigma^j - \rho^j \otimes \rho^j) \quad . \quad (6.2)$$

Although it has the same form as (3.3), its content is quite different. With tetrads $e^j = \sigma^j$, $e^{j+3} = \rho^j$ ($j = 1, 2, 3$) curvature 2-forms are given by

$$\mathcal{R}_{j,k} = -\mathcal{R}_{j+3,k+3} = \frac{1}{4}(e^j \wedge e^k - e^{j+3} \wedge e^{k+3})$$

$$\mathcal{R}_{j,k+3} = -\frac{1}{4}(e^j \wedge e^{k+3} - e^k \wedge e^{j+3}) \quad , \quad (j, k = 1, 2, 3) \quad . \quad (6.3)$$

Ricci tensors are

$$(Ricci)_{ab} = \eta_{ab} = diag(1, 1, 1, -1, -1, -1) \quad . \quad (6.4)$$

The manifold $SO(3, 1)$ is maximally symmetric. Its scalar curvature is non-vanishing.

The manifold is not compact. There may be configurations with x -dependent Φ which solve the equations in (1.2). The unboundedness of the manifold make it possible to have such configurations, though we have not so far found one. In Weyl invariant spacetime the Kalb-Ramond field strengths $H \sim H_\alpha$ or H_β where

$$\begin{aligned} H_\alpha &= +e^1 \wedge e^2 \wedge e^3 - e^1 \wedge e^5 \wedge e^6 + e^2 \wedge e^4 \wedge e^6 - e^3 \wedge e^4 \wedge e^5 \quad , \\ H_\beta &= -e^4 \wedge e^5 \wedge e^6 + e^2 \wedge e^3 \wedge e^4 - e^1 \wedge e^3 \wedge e^5 + e^1 \wedge e^2 \wedge e^6 \end{aligned} \quad (6.5)$$

would play an important role. H_α and H_β satisfy

$$\begin{aligned} dH_\alpha &= dH_\beta = 0 \quad , \\ *H_\alpha &= H_\beta \quad , \quad *H_\beta = H_\alpha \quad , \\ (S_\alpha)_{ab} &= +\eta_{ab} \quad , \quad (S_\beta)_{ab} = -\eta_{ab} \quad . \end{aligned} \quad (6.6)$$

7. Discussions

In this paper we have investigated Weyl invariant spacetime defined as a background solving the set of the equations in (1.2). Beside the well-known Ricci flat spacetime we have found a new class of backgrounds with nonvanishing Ricci tensors. The Kalb-Ramond field strengths assume nontrivial configurations, balancing the two sides of the Einstein equations. The scalar curvature vanishes as the dilaton field is set to be constant.

The construction of Weyl invariant spacetime is based on the fact that the Kalb-Ramond field strengths H are 3-form fields satisfying $dH = *d(*H) = 0$. We prepare a six-dimensional manifold in the form of a product of two three-dimensional group manifolds. On each three-dimensional group manifold, $H = e^1 \wedge e^2 \wedge e^3$ where e^j 's are

Maurer-Cartan forms solves the equations for H . We need two group manifolds so as to have vanishing scalar curvature in six dimensions. One of the two group manifolds must have negative scalar curvature, which is achieved either by preparing a group manifold with intrinsically negative curvature or by flipping the signature in the metric.

In either case there have appeared timelike dimensions which makes it difficult to apply our results to, say, the string theory. Excitations in timelike dimensions would cause instability. It has to be explored if there is a way to control excitations in timelike dimensions. Many arguments have been put forward in the literature that extra timelike dimensions can physically make sense when certain conditions are met.[9]-[12] Another related point to be clarified is about the existence of supersymmetry on the Weyl invariant manifolds.

In the examples discussed above the Kalb-Ramond field strengths have constant magnitude. They are of the monopole type in the sense that they cannot be written globally as $H = dB$. What would be dynamics of strings on such backgrounds? It is about strings on group manifolds.[13] Motion of particles in a constant magnetic field $F = dA$ or of open strings in a constant B effectively produces non-commutative geometry. What would be dynamics of closed strings in a constant H on a group manifold?

The first example of the Weyl invariant spacetime, $SU(2) \otimes \overline{SU(2)}$, has many interesting properties. First of all the manifold is compact. Secondly there is one parameter family of Weyl invariant spacetime on this manifold. H is given by (3.11). It is interesting to know the relevance of the parameter β in (3.11). The configuration H_0 in (3.7) bridges the two $SU(2)$ manifolds. Thirdly there is accidental symmetry of the manifold, namely symmetry under the interchange of the two $SU(2)$'s. Fourthly a six-dimensional flat spacetime with three timelike dimensions has Majorana-Weyl spinors.

Motivated by the string theory, we have investigated Weyl invariant spacetime in general framework. It would be of great interest to know if the Weyl invariant spacetime discussed in the present paper has applications in physics. We hope to come back to this point in near future.

Acknowledgments

I benefited from enlightening discussions with many of my colleagues including K. Higashijima, T. Nakatsu, T. Ohnishi, N. Ohta, M. Sato, T. Tanaka, and K. Tsuda. I would like to duly thank them. This work was supported in part by Scientific Grants from the Ministry of Education and Science, Grant No. 13135215 and Grant No. 13640284.

References

- [1] C.G. Callan, D. Friedan, E.J. Martinec and M.J. Perry, *Nucl. Phys.* **B262** (1985) 593.
- [2] E. Braaten, T.L. Curtright and C.K. Zachos, *Nucl. Phys.* **B260** (1985) 630.
- [3] M. Kalb and P. Ramond, *Phys. Rev.* **D9** (1974) 2273.
- [4] Y. Hosotani, *Phys. Lett.* **B147** (1984) 44.
- [5] A. Hosoya and W. Ogura, *Phys. Lett.* **B225** (1989) 117;
S.J. Rey, *Nucl. Phys.* **B336** (1990) 146.
- [6] H. Emoto, Y. Hosotani and T. Kubota, *Prog. Theoret. Phys.* **108** (2002) 157 [hep-th/0201141];
Y. Hosotani, H. Emoto and T. Kubota, hep-ph/0209112.
- [7] P. Orland, *Nucl. Phys.* **B205** (1982) 107.
- [8] P.A.M. Dirac, *Proc. Roy. Soc.* **A133** (1931) 60;
T.T. Wu and C.N. Yang, *Phys. Rev.* **D12** (1975) 3845.
- [9] A.D. Sakharov, *Zh. Eksp. Teor. Fiz.* **87** (1984) 375 [*Sov. Phys. JETP* **60** (1984) 214];
I.Ya. Aref'eva and I.V. Volvovich, *Phys. Lett.* **B164** (1985) 287;
I.Ya. Aref'eva, B.G. Dragovich and I.V. Volvovich, *Phys. Lett.* **B177** (1986) 357;
A. Chamblin and R. Emparan, *Phys. Rev.* **D55** (1997) 754;
M. Chaichian and A.B. Kobakhidze, *Phys. Lett.* **B488** (2000) 117.
- [10] C. Vafa, *Nucl. Phys.* **B496** (1996) 403.
- [11] F.J. Yndurain, *Phys. Lett.* **B256** (1991) 15;
G. Dvali, G. Gabadadze, and G. Senjanovic, hep-ph/9910207.
- [12] S. Matsuda and S. Seki, *Phys. Rev.* **D63** (2001) 065014; *Nucl. Phys.* **B599** (2001) 119.
- [13] D. Gepner and E. Witten, *Nucl. Phys.* **B278** (1986) 493;
J. Maldacena and H. Ooguri, *J. Math. Phys.* **42** (2001) 2929;
J. Maldacena, H. Ooguri, and J. Son, *J. Math. Phys.* **42** (2001) 2961.